○イヌタデとハナタデの雑種（檜山庫三）　Kōzō HIYAMA: A new hybrid of *Persicaria*

1956年10月4日，東京都練馬区の武蔵大学構内で変つたタデの1種を見つけた。全形は一見したところハナタデ（ヤブタデ）に似たもので，茎は緑色で立ち，節間は1-3 cmばかりで短く，葉はおおむね長楕円状披針形，漸尖頭ではあるが，先の方が尾状となるようなことがなく，葉脚も尖り，長は6-9 cm，巾は2-2.5 cmで，質が薄く，葉面に毛はあるがごく少く，表面は中脈上と葉縁に近い部分に短毛（ハナタデのように長くない）がまばらに生え，また裏面の脈上にも多少の疎毛があり，葉縁毛もハナタデのものほど長くはない。花部や果実はハナタデと余り変らない。このタデは上記の如く，葉形はイヌタデ（ハナタデ）に似ているが，それよりも巾が広く，葉質はハナタデのようである。その生えていた場所も，イタタデとハナタデとが接して生えていた所であつたから，おそらくは両者の間の自然雑種ではないかと考えられる。そこで，この者にイヌハナタデ（*Persicaria* × *musashinoensis* Hiyama）の名を与えておきたいと思う。

Persicaria × musashinoensis Hiyama, hybr. nov.

Persicaria longiseta (De Bruyn) Kitagawa × *P. Yokusaiana* (Makino) Nakai.

Persicaria Yokusaianae affinis sed ab ea foliorum forma differt.

Caulis erectiusculus 50 cm altus ramosus, ramis subdivaricato-ascendentibus, viridescens glaber, internodii 1-3 cm longi. Ochreae pallescentes hyalino-membranaceae nervatae parce hirsutae 4-8 mm longae apice longe setoso-ciliatae. Folia infima ovato-elliptica minora, mediana oblongo-lanceolata apicem versus sensim attenuata basi acuta, textu submembranacea, 6-5 cm longa 2-2.5 cm lata supra ad nervum medianum et prope marginem parce adpresse hirsuta subtus ad nervum tantum parce pubigera, utrinque nervis lateralibus 10-12 supra faciem subimpressis. Flores et fructus fere ut *P. Yokusaiana*.—Nom. Jap. Inu-hanatade, nov.

Hab. Hondo: Egoda, Nerima-ku, Tokyo, Prov. Musashi (Hiyama—Oct. 4, 1956—typus in Herb. Nat. Mus. Tokyo).

This plant is probably a hybrid between *Persicaria longisetsa* and *P. Yokusaiana*. It may be distinguished from the former chiefly by its thinner texture of leaves and its looser inflorescence. ochreas are sparsely pubescent and fringed at the top with slender bristles (4-8 mm long). Flowers are arranged in rather loose spike-like racemes, somewhat interrupted below, pinkish, appearing in autumn. Calyx about 2 mm long, 5-parted. Styles 3. Achene 3-gonous, 1.5-2 mm long, blackish brown, lucid.